

## Indications

Constipation and its accessory symptoms such as heavy head, hot flashes, eczema/dermatitis, pimple (acne), loss of appetite (anorexia), abdominal bloating, abnormal intestinal fermentation, and hemorrhoid.

<Caution related to indications>

This medicine can be used regardless of the physical strength.



## ! Caution for use

### To be avoided

1. Don't give this medicine to children under 3 months of age.
2. This medicine should not be taken together with the following medicines:  
Other cathartic agents.
3. Women who are nursing should not be taken this medicine. If they take it, avoid breast feeding.

### Who should seek consultation

1. The following persons should consult a physician, pharmacist or registered distributor before taking this medicine.
  - (1) Persons undergoing medical treatment from a physician.
  - (2) Pregnant women or women suspected of being pregnant.
  - (3) Persons who is physically weak (declining of physical strength or constitutional weakness).
  - (4) Persons with weak stomach and intestine who have a tendency to diarrhea.
  - (5) Elderly persons.
  - (6) Persons with the following symptom:  
Edema.
  - (7) Persons diagnosed as having the following:  
Hypertension, heart disease or kidney disease.
2. The following symptoms may be adverse reactions of this medicine. If any of the following symptoms occur, stop taking this medicine immediately and consult a physician, pharmacist or registered distributor with this instruction leaflet.

Areas	Symptoms
Gastrointestinal system	Diarrhea with severe abdominal pain, abdominal pain

In very rare cases the following serious symptoms may occur. In this case, consult a physician immediately.

Symptoms' name	Symptoms
Pseudoaldosteronism, myopathy	In addition to listlessness, numbness, stretched feeling, and stiffness of hands and feet, feeling of weakness and muscular pain will appear and gradually progress

3. A following symptom may appear after taking this medicine. If the symptom continues or worsen, consult a physician, pharmacist, or registered distributor with this leaflet:  
Diarrhea.
4. Stop taking this medicine and consult a physician, pharmacist or registered distributor with this leaflet, if symptoms do not improve after taking this medicine for 5 to 6 days.
5. If you take this medicine regularly for a long period, consult a physician pharmacist or registered distributor.

## Components

Daiokanzoto (大黄甘草湯) is extract from mixture of the following crude drugs;  
Rhubarb (大黄), Glycyrrhiza (甘草)

\*If you have any questions, please consult to a physician, pharmacist, or registered distributor .  
Supervised by Ichiro Arai Ph.D., Professor, Division of Kampo Pharmaceutical Sciences, Nihon Pharmaceutical University

