

Indications

Following symptoms in persons with substantial physical strength who suffer from flank to solar plexus and have a tendency of constipation:

Gastritis, habitual constipation, and accessory symptoms of hypertension or obesity (stiff shoulder, headache, constipation, neurosis and obesity).

! Caution for use

To be avoided

1. Don't give this medicine to children under 3 months of age.
2. This medicine should not be taken together with the following medicines:
Other cathartic agents.
3. Women who are nursing should not be taken this medicine. If they take it, avoid breast feeding.

Who should seek consultation

1. The following persons should consult a physician, pharmacist or registered distributor before taking this medicine.
 - (1) Persons undergoing medical treatment from a physician.
 - (2) Pregnant women or women suspected of being pregnant.
 - (3) Persons who is physically weak (declining of physical strength or constitutional weakness).
 - (4) Persons with weak stomach and intestine who have a tendency to diarrhea.
 - (5) Persons who have experienced rash, redness, itching, etc. due to medicines, etc.
2. The following symptoms may be adverse reactions of this medicine. If any of the following symptoms occur, stop taking this medicine immediately and consult a physician, pharmacist or registered distributor with this instruction leaflet.

Areas	Symptoms
Skin	Rash, redness, itching
Gastrointestinal system	Diarrhea with severe abdominal pain, abdominal pain

In very rare cases the following serious symptoms may occur. In this case, consult a physician immediately.

Symptoms' name	Symptoms
Interstitial pneumonia	Shortness and / or difficulties of breath when go upstairs or overwork, sudden dry cough and / or fever and its continuance.
Hepatic function failure	The following symptoms appear: Fever, itching, rash, jaundice (yellowing of skin and white of eyes), brown urine, whole-body tiredness, loss of appetite

3. A following symptom may appear after talking this medicine. If the symptom continues or worsen, consult a physician, pharmacist, or registered distributor with this leaflet :
Diarrhea.
4. Stop taking this medicine and consult a physician, pharmacist or registered distributor with this leaflet, if symptoms do not improve after taking this medicine for about 1 month (5 to 6 days if you take this medicine for habitual constipation and constipation associated with hypertension or obesity).

Components

Daisaikoto (大柴胡湯) is extract from mixture of the following crude drugs;
Bupleurum Root (柴胡), Pinellia Tuber (半夏), Ginger (生姜), Scutellaria Root (黄芩), Peony Root (芍薬),
Jujube (大枣), Immature Orange (枳实), Rhubarb (大黄) .

*If you have any questions, please consult to a physician, pharmacist, or registered distributor .
Supervised by Ichiro Arai Ph.D., Professor, Division of Kampo Pharmaceutical Sciences, Nihon Pharmaceutical University