

Indications

Following symptoms in persons with moderate or more physical strength:
Frequent urination, micturition pain, feeling of residual urine and cloudy urine.

! Caution for use

To be avoided

Don't give this medicine to children under 3 months of age.

Who should seek consultation

- The following persons should consult a physician, pharmacist or registered distributor before taking this medicine.
 - Persons undergoing medical treatment from a physician.
 - Pregnant women or women suspected of being pregnant.
 - Persons with weak stomach and intestine who have a tendency to diarrhea.
 - Elderly persons.
 - Persons with the following symptom:
 - Edema.
 - Persons diagnosed as having the following:
 - Hypertension, heart disease or kidney disease.
- The following symptoms may be adverse reactions of this medicine. If any of the following symptoms occur, stop taking this medicine immediately and consult a physician, pharmacist or registered distributor with this instruction leaflet.

Areas	Symptoms
Gastrointestinal system	Loss of appetite, gastric distress

In very rare cases the following serious symptoms may occur. In this case, consult a physician immediately.

Symptoms' name	Symptoms
Interstitial pneumonia	Shortness and / or difficulties of breath when go upstairs or overwork, sudden dry cough and/or fever and its continuance
Pseudoaldosteronism, myopathy	In addition to listlessness, numbness, stretched feeling, and stiffness of hands and feet, feeling of weakness and muscular pain will appear and gradually progress
Mesenteric phlebosclerosis	Long-term use of this medicine may cause repetitive appearance of following symptoms: Abdominal pain, diarrhea, constipation, abdominal bloating

- A following symptom may appear after taking this medicine. If the symptom continues or worsen, consult a physician, pharmacist, or registered distributor with this leaflet:
 - Diarrhea.
- Stop taking this medicine and consult a physician, pharmacist or registered distributor with this leaflet if symptoms do not improve after taking this medicine for about 1 month.
- If you take this medicine regularly for a long period, consult a physician, pharmacist, or registered distributor.

Components

Gorinsan (五淋散) is extract from mixture of the following crude drugs;
Poria Sclerotium (茯苓), Japanese Angelica Root (当归), Scutellaria Root (黄芩), Glycyrrhiza (甘草), Peony Root (芍药), Gardenia Fruit (山梔子), Rehmannia Root (地黄), Alisma Tuber (沢瀉), Akebia Stem (木通), Aluminum Silicate Hydrate with Silicon Dioxide (滑石), Plantago Seed (車前子).
(Products without Rehmannia Root, Alisma Tuber, Akebia Stem, Aluminum Silicate Hydrate with Silicon Dioxide and Plantago Seed can be possible.)

*If you have any questions, please consult to a physician, pharmacist, or registered distributor.
Supervised by Ichiro Arai Ph.D., Professor, Division of Kampo Pharmaceutical Sciences, Nihon Pharmaceutical University

