

Indications

Following symptoms in persons with moderate or poor physical strength who tend to get tired easily, tend to be sensitive to cold in hands and feet, have decreased or increased urine volume, and/or sometimes have dry mouth:

Leg pain, lower back pain, numbness, blurred eye in elderly, itching, difficulty in urination, residual urine, nocturia, frequent urination, edema, associated symptoms of hypertension (stiff shoulder, heavy head, ear noise), and mild incontinence.

! Caution for use

To be avoided

This drug is not to be taken:

- (1) Children under 3 months of age.
- (2) Persons with weak stomach and intestine.
- (3) Persons who have a tendency to diarrhea.

Who should seek consultation

1. The following persons should consult a physician, pharmacist or registered distributor before taking this medicine.
 - (1) Persons undergoing medical treatment from a physician.
 - (2) Pregnant women or women suspected of being pregnant.
 - (3) Red faced persons with hot flashes who have fulfilling physical performance.
 - (4) Persons who have experienced rash, redness, itching, etc. due to medicines or something.
2. The following symptoms may be adverse reactions of this medicine. If any of the following symptoms occur, stop taking this medicine immediately and consult a physician, pharmacist, or registered distributor with this instruction leaflet.

Areas	Symptoms
Skin	Rash, redness, itching
Gastrointestinal system	Loss of appetite, gastric distress, gastrointestinal pain
Others	Palpitations, hot flush, numbness of lip or tongue

3. A following symptom may appear after taking this medicine. If the symptom continues or worsen, consult a physician, pharmacist, or registered distributor with this leaflet:
Diarrhea.
4. Stop taking this medicine and consult a physician, pharmacist or registered distributor with this leaflet, if symptoms do not improve after taking this medicine for about 1 month.

Components

Hachimijogon (八味地黄丸) is extract from mixture of the following crude drugs;

Rehmannia Root (地黄), Cornus Fruit (山茱萸), Dioscorea Rhizome (山药), Alisma Rhizome (沢瀉), Poria Sclerotium (茯苓), Moutan Bark (牡丹皮), Cinnamon Bark (桂皮), Processed Aconite Root (加工ブシ).

*If you have any questions, please consult to a physician, pharmacist, or registered distributor .
Supervised by Ichiro Arai Ph.D., Professor, Division of Kampo Pharmaceutical Sciences, Nihon Pharmaceutical University