

# HANGEKOBOKUTO (半夏厚朴湯)

## Indications

Following symptoms in person with moderate physical strength who feel depressed, experience feeling of a foreign body in the throat and esophagus, and sometimes experience palpitations, dizziness, nausea, etc:

Choking feeling in the throat, nervous gastritis, anxiety, cough, hoarseness, and morning sickness.

## ! Caution for use

### To be avoided

Don't give this medicine to children under 3 months of age.

### Who should seek consultation

- The following persons should consult a physician, pharmacist or registered distributor before taking this medicine.
  - Persons undergoing medical treatment from a physician.
  - Persons who have experienced rash, redness, itching, etc. due to medicines, etc.
- The following symptoms may be adverse reactions of this medicine. If any of the following symptoms occur, stop taking this medicine immediately and consult a physician, pharmacist or registered distributor with this instruction leaflet.

Areas	Symptoms
Skin	Rash, redness, itching

- Stop taking this medicine and consult a physician, pharmacist or registered distributor with this leaflet, if symptoms do not improve after taking this medicine for about 1 month (about 1 week if you take this medicine for morning sickness).

## Components

Hangekobokuto is extract from mixture of the following crude drugs;

Pinellia Tuber (半夏), Poria Sclerotium (茯苓), Magnolia Bark (厚朴), Ginger (生姜), Perilla Herb (蘇葉).

\*If you have any questions, please consult to a physician, pharmacist, or registered distributor .  
Supervised by Ichiro Arai Ph.D., Professor, Division of Kampo Pharmaceutical Sciences, Nihon Pharmaceutical University