

## Indications

Following symptoms in person with moderate or more physical strength:

Initial stage of common cold (without spontaneous sweating), coryza, rhinitis, headache, shoulder stiffness, muscle pain and pain in hands and shoulder.

## ! Caution for use

### To be avoided

Don't give this medicine to children under 3 months of age.

### Who should seek consultation

- The following persons should consult a physician, pharmacist or registered distributor before taking this medicine.
  - Persons undergoing medical treatment from a physician.
  - Pregnant women or women suspected of being pregnant.
  - Persons who is physically weak (declining of physical strength or constitutional weakness).
  - Persons who have gastrointestinal weakness.
  - Persons with excessive sweating.
  - Elderly persons.
  - Persons who have experienced rash, redness, itching, etc. due to medicines, etc.
  - Persons with the following symptoms:  
Edema or urination difficulty.
  - Persons diagnosed as having the following:  
Hypertension, heart disease, kidney disease or thyroid disease.
- The following symptoms may be adverse reactions of this medicine. If any of the following symptoms occur, stop taking this medicine immediately and consult a physician, pharmacist or registered distributor with this instruction leaflet.

Areas	Symptoms
Skin	Rash, redness, itching
Gastrointestinal system	Nausea, loss of appetite, gastric distress

In very rare cases the following serious symptoms may occur. In this case, consult a physician immediately.

Symptoms' name	Symptoms
Pseudoaldosteronism, myopathy	In addition to listlessness, numbness, stretched feeling, and stiffness of hands and feet, feeling of weakness and muscular pain will appear and gradually progress
Hepatic function failure	The following symptoms appear: Fever, itching, rash, jaundice (yellowing of skin and white of eyes), brown urine, whole-body tiredness, loss of appetite

- Stop taking this medicine and consult a physician, pharmacist or registered distributor with this leaflet, if symptoms do not improve after taking this medicine for about 1 month (5 to 6 times if you take this medicine for the initial stage of the common cold, coryza and headache).
- If you take this medicine regularly for a long period, consult a physician, pharmacist, or registered distributor.

## Components

Kakkonto (葛根湯) is extract from the mixture of the following crude drugs;

Pueraria Root (葛根), Ephedra Herb (麻黄), Jujube (大枣), Cinnamon Bark (桂皮), Peony Root (芍药), Glycyrrhiza (甘草), Ginger (生姜).



\*If you have any questions, please consult to a physician, pharmacist, or registered distributor .

Supervised by Ichiro Arai Ph.D., Professor, Division of Kampo Pharmaceutical Sciences, Nihon Pharmaceutical University