

Indications

Following symptoms in persons with moderate or poor physical strength who have mental and physical tiredness and pale skin, and have feeling of warmth in some case:

Anemia, insomnia, mental anxiety, and neurosis.

! Caution for use

To be avoided

Don't give this medicine to children under 3 months of age.

Who should seek consultation

1. The following persons should consult a physician, pharmacist or registered distributor before taking this medicine.

- (1) Persons undergoing medical treatment from a physician.
- (2) Pregnant women or women suspected of being pregnant.
- (3) Elderly persons.
- (4) Persons who have experienced rash, redness, itching, etc. due to medicines, etc.
- (5) Persons with the following symptom:
Edema.
- (6) Persons diagnosed as having the following:
Hypertension, heart disease or kidney disease.

2. The following symptoms may be adverse reactions of this medicine. If any of the following symptoms occur, stop taking this medicine immediately and consult a physician, pharmacist or registered distributor with this instruction leaflet.

Areas	Symptoms
Skin	Rash, redness, itching

In very rare cases the following serious symptoms may occur. In this case, consult a physician immediately.

Symptoms' name	Symptoms
Pseudoaldosteronism, myopathy	In addition to listlessness, numbness, stretched feeling, and stiffness of hands and feet, feeling of weakness and muscular pain will appear and gradually progress
Mesenteric phlebosclerosis	Long-term use of this medicine may cause repetitive appearance of following symptoms: Abdominal pain, diarrhea, constipation, abdominal bloating

3. Stop taking this medicine and consult a physician, pharmacist, or registered distributor with this leaflet, if symptoms do not improve after taking this medicine for about 1 month.
4. If you take this medicine regularly for a long period, consult a physician, pharmacist, or registered distributor.

Components

Kamikihito (加味帰脾湯) is extract from mixture of the following crude drugs;
Ginseng (人參), Atractylodes Rhizome (白朮) or Atractylodes Lancea Rhizome (蒼朮), Poria Sclerotium (茯苓), Jujube seed (酸棗仁), Longan Aril (龍眼肉), Astragalus Root (黃耆), Japanese Angelica Root (當歸), Polygala Root (遠志), Bupleurum root (柴胡), Gardenia Fruit (山梔子), Glycyrrhiza (甘草), Saussurea Root (木香), Jujube (大棗), Ginger (生姜), (Moutan Bark (牡丹皮)).

*If you have any questions, please consult to a physician, pharmacist, or registered distributor.
Supervised by Ichiro Arai Ph.D., Professor, Division of Kampo Pharmaceutical Sciences, Nihon Pharmaceutical University

