

## Indications

Following symptoms in red-faced persons with moderate or more physical strength who get hot flashes easily and become irritated and restless easily:

Nosebleed, insomnia, neurosis, gastritis, hangover, chi-no-michi-sho\*, dizziness, palpitation, menopausal disorders, eczema/dermatitis, itching, and stomatitis.

\*Chi-no-michi-sho is a generic term of neuropsychiatric symptoms such as anxiety and irritation and physical symptoms due to fluctuation of female hormones caused by menstruation, pregnancy, childbirth or menopause.

## ! Caution for use

### To be avoided

Don't give this medicine to children under 3 months of age.

### Who should seek consultation

- The following persons should consult a physician, pharmacist or registered distributor before taking this medicine.
  - Persons undergoing medical treatment from a physician.
  - Pregnant women or women suspected of being pregnant.
  - Persons who is physically weak (declining of physical strength or constitutional weakness).
- The following symptoms may be adverse reactions of this medicine. In very rare cases, the following serious symptoms may occur. In this case, stop taking this medicine and consult a physician with this instruction leaflet immediately.

Symptoms' name	Symptoms
Interstitial pneumonia	Shortness and/or difficulties of breath when go upstairs or overwork, sudden dry cough and / or fever and its continuance
Hepatic function failure	The following symptoms appear: Fever, itching, rash, jaundice (yellowing of skin and white of eyes), brown urine, whole-body tiredness, loss of appetite
Mesenteric phleboscrosis	Long-term use of this medicine may cause repetitive appearance of following symptoms: Abdominal pain, diarrhea, constipation, abdominal bloating

- Stop taking this medicine and consult a physician, pharmacist or registered distributor with this leaflet, if symptoms do not improve after taking this medicine for about 1 month (5 to 6 days if you take this medicine for nosebleed or hangover).
- If you take this medicine regularly for a long period, consult a physician, pharmacist, or registered distributor.

## Components

Orengedokuto (黄連解毒湯) is extract from mixture of the following crude drugs;  
Coptis Rhizome (黄連), Scutellaria Root (黄芩), Phellodendron Bark (黄柏), Gardenia Fruit (山梔子).

\*If you have any questions, please consult to a physician, pharmacist, or registered distributor .  
Supervised by Ichiro Arai Ph.D., Professor, Division of Kampo Pharmaceutical Sciences, Nihon Pharmaceutical University