

## Indications

Following symptoms in persons with moderate and poor physical strength who cannot clear up throat easily, and sometimes experience violent coughing or dryness of the throat:

Dry cough, bronchitis, bronchial asthma, and pharyngitis, croak (hoarseness).

## ! Caution for use

### To be avoided

Don't give this medicine to children under 3 months of age.

### Who should seek consultation

1. The following persons should consult a physician, pharmacist, or registered distributor before taking this medicine.

- (1) Persons undergoing medical treatment from a physician.
- (2) Pregnant women or women suspected of being pregnant.
- (3) Persons expelling large amount of watery phlegm.
- (4) Elderly persons.
- (5) Persons with the following symptoms:

Edema.

(6) Persons diagnosed as having the following:

Hypertension, heart disease, kidney disease or thyroid disease.

2. The following symptoms may be adverse reactions of this medicine. If any of the following symptoms occur, stop taking this medicine immediately and consult a physician, pharmacist, or registered distributor with this instruction leaflet.

Areas	Symptoms
Gastrointestinal system	Loss of appetite, gastric distress

In very rare cases the following serious symptoms may occur. In this case, consult a physician immediately.

Symptoms' name	Symptoms
Interstitial pneumonia	Shortness and/or difficulties of breath when go upstairs or overwork, sudden dry cough and/or fever and its continuance
Pseudoaldosteronism, myopathy	In addition to listlessness, numbness, stretched feeling, and stiffness of hands and feet, feeling of weakness and muscular pain will appear and gradually progress
Hepatic function failure	The following symptoms appear: Fever, itching, rash, jaundice (yellowing of skin and white of eyes), brown urine, whole-body tiredness, loss of appetite

3. Stop taking this medicine and consult a physician, pharmacist or registered distributor with this leaflet, if symptoms do not improve after taking this medicine for about 1 month (about 1 week if you take this medicine for dry cough).
4. If you take this medicine regularly for a long period, consult a physician, pharmacist or registered distributor.

## Components

Bakumondoto (麦門冬湯) is extract from mixture of the following crude drugs;

Ophiopogon Tuber (麦門冬), Brown Rice (粳米), Ginseng (人參), Pinellia Tuber (半夏), Jujube (大棗), Glycyrrhiza (甘草).



\*If you have any questions, please consult to a physician, pharmacist, or registered distributor .

Supervised by Ichiro Arai Ph.D., Professor, Division of Kampo Pharmaceutical Sciences, Nihon Pharmaceutical University

