

## Indications

Following symptoms in persons who complaint thirst and decreased urine volume accompanying any of dizziness, nausea, vomiting, abdominal pain, headache or edema. This medicine can be applied to everyone regardless of physical strength:

Watery diarrhea, acute gastroenteritis (Do not use for **shiburihara** [tenesmus]\*), heat stroke, headache, edema, and hangover.

\***Shiburihara** [tenesmus] is the symptom consisting of feeling of unsatisfied defecation and repetitive defecation desire with abdominal pain.

## ! Caution for use

### To be avoided

Don't give this medicine to children under 3 months of age.

### Who should seek consultation

- The following persons should consult a physician, pharmacist or registered distributor before taking this medicine.
  - Persons undergoing medical treatment from a physician.
  - Pregnant women or women suspected of being pregnant.
  - Persons who have experienced rash, redness, itching, etc. due to medicines, etc.
- The following symptoms may be adverse reactions of this medicine. If any of the following symptoms occur, stop taking this medicine immediately and consult a physician, pharmacist or registered distributor with this instruction leaflet.

Areas	Symptoms
Skin	Rash, redness, itching

- Stop taking this medicine and consult a physician, pharmacist or registered distributor with this leaflet, if symptoms do not improve after taking this medicine for about 1 month (5 to 6 times if you take this medicine for acute gastroenteritis or hangover, and 5 to 6 days if you take it for watery diarrhea or heat stroke).

## Components

Goreisan (五苓散) is extract from mixture of the following crude drugs;

Alisma Tuber (沢瀉), Polyporus Sclerotium (猪苓), Poria Sclerotium (茯苓), Atractylodes Lancea Rhizome (蒼朮) or Atractylodes Rhizome (白朮), Cinnamon Bark (桂皮) .