

Indications

Following symptoms in persons with moderate or more physical strength:
Nasal stuffiness, sinus problem, and chronic rhinitis.

! Caution for use

To be avoided

Don't give this medicine to children under 3 months of age.

Who should seek consultation

- The following persons should consult a physician, pharmacist or registered distributor before taking this medicine.
 - (1) Persons undergoing medical treatment from a physician.
 - (2) Pregnant women or women suspected of being pregnant.
 - (3) Persons who is physically weak (declining of physical strength or constitutional weakness).
 - (4) Persons who have gastrointestinal weakness.
 - (5) Persons with excessive sweating.
 - (6) Elderly persons.
 - (7) Persons who have experienced rash, redness, itching, etc. due to medicines or something.
 - (8) Persons with the following symptoms:
Edema, difficulty or urination difficulty.
 - (9) Persons diagnosed as having the following:
Hypertension, heart disease, kidney disease, or thyroid disease.
- The following symptoms may be adverse reactions of this medicine. If any of the following symptoms occur, stop taking this medicine immediately and consult a physician, pharmacist or registered distributor with this instruction leaflet.

Areas	Symptoms
Skin	Rash, redness, itching
Gastrointestinal system	Nausea, loss of appetite, gastric distress

In very rare cases the following serious symptoms may occur. In this case, consult a physician immediately.

Symptoms' name	Symptoms
Pseudoaldosteronism, myopathy	In addition to listlessness, numbness, stretched feeling, and stiffness of hands and feet, feeling of weakness and muscular pain will appear and gradually progress

- Stop taking this medicine and consult a physician, pharmacist or registered distributor with this leaflet, if symptoms do not improve after taking this medicine for about 1 month.
- If you take this medicine regularly for a long period, consult a physician, pharmacist or registered distributor.

Components

Kakkontokasenkyushin' i (葛根湯加川芎辛夷) is extract from mixture of the following crude drugs;
Pueraria Root (葛根), Jujube (大棗), Ephedra Herb (麻黃), Glycyrrhiza (甘草), Cinnamon Bark (桂皮),
Peony Root (芍藥), Magnolia Flower (辛夷), Cnidium Rhizome (川芎), Ginger (生姜).



*If you have any questions, please consult to a physician, pharmacist, or registered distributor .

Supervised by Ichiro Arai Ph.D., Professor, Division of Kampo Pharmaceutical Sciences, Nihon Pharmaceutical University

