

Indications

This medicine can be used regardless of physical strength and is indicated to the following symptoms with pain associated with sudden muscle spasm:

Cramps in the calves, muscle spasm, abdominal pain, low back pain.

! Caution for use

To be avoided

1. Don't give this medicine to children under 3 months of age.
2. The following persons should not take this medicine:
Persons diagnosed as having the following:
Heart disease.
3. This medicine should only be taken when you have a symptom, and avoid taking it for a long time.

Who should seek consultation

1. The following persons should consult a physician, pharmacist, or registered distributor before taking this medicine.
 - (1) Persons undergoing medical treatment from a physician.
 - (2) Pregnant women or women suspected of being pregnant.
 - (3) Elderly persons.
 - (4) Persons with the following symptoms:
Edema.
 - (5) Persons diagnosed as having the following:
Hypertension or kidney disease.
2. The following symptoms may be adverse reactions of this medicine. If any of the following symptoms occur, stop taking this medicine immediately and consult a physician, pharmacist or registered distributor with this instruction leaflet. In very rare cases the following serious symptoms may occur. In this case, consult a physician immediately.

Symptoms' name	Symptoms
Congestive heart failure, ventricular tachycardia	Whole body tiredness, palpitations, shortness of breath, chest discomfort, chest pain, dizziness and faint may appear
Hepatic function failure	The following symptoms appear: Fever, itching, rash, jaundice (yellowing of skin and white of eyes), brown urine, whole-body tiredness, loss of appetite
Interstitial pneumonia	Shortness and/or difficulties of breath when go upstairs or overwork, sudden dry cough and/or fever and its continuance
Pseudoaldosteronism, myopathy	In addition to listlessness, numbness, stretched feeling, and stiffness of hands and feet, feeling of weakness and muscular pain will appear and gradually progress

3. Stop taking this medicine and consult a physician, pharmacist or registered distributor with this leaflet, if symptoms do not improve after taking this medicine 5 to 6 times.

Components

Shakuyakukanzoto (芍薬甘草湯) is extract from mixture of the following crude drugs;
Peony Root (芍薬), Glycyrrhiza (甘草).

*If you have any questions, please consult to a physician, pharmacist, or registered distributor.
Supervised by Ichiro Arai Ph.D., Professor, Division of Kampo Pharmaceutical Sciences, Nihon Pharmaceutical University