

Indications

Following symptoms in persons with moderate or slightly poor physical strength who suffer from coughing with thin watery phlegm and runny nose:

Bronchitis, bronchial asthma, rhinitis, allergic rhinitis, edema, common cold, and hay fever.

! Caution for use

To be avoided

Don't give this medicine to children under 3 months of age.



Who should seek consultation

1. The following persons should consult a physician, pharmacist or registered distributor before taking this medicine.

- (1) Persons undergoing medical treatment from a physician.
- (2) Pregnant women or women suspected of being pregnant.
- (3) Persons who is physically weak (declining of physical strength or constitutional weakness).
- (4) Persons who have gastrointestinal weakness.
- (5) Persons with excessive sweating.
- (6) Elderly persons.
- (7) Persons who have experienced rash, redness, itching, etc. due to medicines, etc.
- (8) Persons with the following symptoms:
Edema or urination difficulty .

- (9) Persons diagnosed as having the following:
Hypertension, heart disease, kidney disease or, thyroid disease.

2. The following symptoms may be adverse reactions of this medicine. If any of the following symptoms occur, stop taking this medicine immediately and consult a physician, pharmacist, or registered distributor with this instruction leaflet.

Areas	Symptoms
Skin	Rash, redness, itching
Gastrointestinal system	Nausea, loss of appetite, gastric distress

In very rare cases the following serious symptoms may occur. In this case, consult a physician immediately.

Symptoms' name	Symptoms
Interstitial pneumonia	Shortness and/or difficulties of breath when go upstairs or overwork, sudden dry cough and/or fever and its continuance
Pseudoaldosteronism, myopathy	In addition to listlessness, numbness, stretched feeling, and stiffness of hands and feet, feeling of weakness and muscular pain will appear and gradually progress
Hepatic function failure	The following symptoms appear: Fever, itching, rash, jaundice (yellowing of skin and white of eyes), brown urine, whole-body tiredness, loss of appetite

3. Stop taking this medicine and consult a physician, pharmacist or registered distributor with this leaflet, if symptoms do not improve after taking this medicine for about 1 month (5 to 6 days if you take this medicine for common cold).

4. If you take this medicine regularly for a long period, consult a physician, pharmacist, or registered distributor.

Components

Shoseiryuto (小青竜湯) is extract from mixture of the following crude drugs;

Ephedra Herb (麻黃) , Peony Root (芍藥) , Processed Ginger (乾姜) , Glycyrrhiza (甘草) , Cinnamon Bark (桂皮) , Asiasarum Root (細辛) , Schisandra Fruit (五味子) , Pinellia Tuber (半夏) .

*If you have any questions, please consult to a physician, pharmacist, or registered distributor .

Supervised by Ichiro Arai Ph.D., Professor, Division of Kampo Pharmaceutical Sciences, Nihon Pharmaceutical University



JAPAN KAMPO MEDICINES MANUFACTURERS ASSOCIATION