

Indications

Following symptoms in persons with moderate physical strength who have pain, and have numbness in some case:
Joint pain, nerve pain, low back pain, and muscle pain.

! Caution for use

To be avoided

Don't give this medicine to children under 3 months of age.

Who should seek consultation

- The following persons should consult a physician, pharmacist or registered distributor before taking this medicine.
 - (1) Persons undergoing medical treatment from a physician.
 - (2) Pregnant women or women suspected of being pregnant.
 - (3) Persons with weak stomach and intestine who have a tendency to diarrhea.
 - (4) Elderly persons.
 - (5) Persons who have experienced rash, redness, itching, etc. due to medicines, etc.
 - (6) Persons with the following symptom:
Edema.
 - (7) Persons diagnosed as having the following:
Hypertension, heart disease or kidney disease.
- The following symptoms may be adverse reactions of this medicine. If any of the following symptoms occur, stop taking this medicine immediately and consult a physician, pharmacist or registered distributor with this instruction leaflet.

Areas	Symptoms
Skin	Rash, redness, itching
Gastrointestinal system	Loss of appetite, gastric distress

In very rare cases the following serious symptoms may occur. In this case, consult a physician immediately.

Symptoms' name	Symptoms
Pseudoaldosteronism, myopathy	In addition to listlessness, numbness, stretched feeling, and stiffness of hands and feet, feeling of weakness and muscular pain will appear and gradually progress

- Stop taking this medicine and consult a physician, pharmacist, or registered distributor with this leaflet, if symptoms do not improve after taking this medicine for about 1 month.
- If you take this medicine regularly for a long period, consult a physician, pharmacist, or registered distributor.

Components

Sokeikakketsuto (疎経活血湯) is extract from mixture of the following crude drugs;
Japanese Angelica Root (当帰), Rehmannia Root (地黄), Cnidium Rhizome (川芎), Atractylodes Lancea Rhizome (蒼朮) (or Atractylodes Rhizome (白朮)), Poria Sclerotium (茯苓), Peach Kernel (桃仁), Peony Root (芍薬),
Achyranthes Root (牛膝), Clematis Root (威靈仙), Sinomenium Stem and Rhizome (防已), Notopterygium (羌活),
Saposhnikovia Root and Rhizome (防風), Japanese Gentian (竜胆), Ginger (生姜), Citrus Unshiu Peel (陳皮),
Angelica Dahurica Root (白芷), Glycyrrhiza (甘草).

*If you have any questions, please consult to a physician, pharmacist, or registered distributor.
Supervised by Ichiro Arai Ph.D., Professor, Division of Kampo Pharmaceutical Sciences, Nihon Pharmaceutical University

